MYCOTAXON

ISSN (print) 0093-4666 (online) 2154-8889 © 2016. Mycotaxon, Ltd.

April–June 2016—Volume 131, pp. 263–268

http://dx.doi.org/10.5248/131.263

Three new species of *Hemicorynespora* and *Solicorynespora* from southern China

Jian Ma[1]*, Ji-Wen Xia[2] & Xiu-Guo Zhang[2]

[1]*College of Agronomy, Jiangxi Agricultural University, Nanchang, 330045, China*

[2]*Department of Plant Pathology, Shandong Agricultural University, Taian, 271018, China*

*Correspondence to: *majian821210@163.com*

Abstract — Three new hyphomycetes were collected from decaying twigs of unidentified plants in southern China. *Hemicorynespora obovoidea* sp. nov., *Solicorynespora jinggangshanensis* sp. nov., and *S. lushanensis* sp. nov. are described, illustrated, and compared with closely related taxa.

Key words — anamorphic fungi, saprobic fungi, taxonomy

Introduction

During ongoing surveys of saprobic microfungi from forests of southern China, several interesting *Corynespora*-like taxa were found on decaying twigs. Among these were three species having the morphological features typical of *Hemicorynespora* M.B. Ellis (Ellis 1972) and *Solicorynespora* R.F. Castañeda & W.B. Kendr. (Castañeda-Ruíz & Kendrick 1990). They differ significantly from previously described species in several morphological features, and therefore are described here as new to science.

Hemicorynespora obovoidea Jian Ma & X.G. Zhang, **sp. nov.** Fig. 1

MycoBank MB 814891

Differs from all other *Hemicorynespora* species by its obovoid, asymmetrically 1-septate conidia.

Type: China, Guangdong Province: Liuxihe National Forest Park, on decaying twigs of an unidentified broadleaf tree, 10 July 2014, J. Ma (**Holotype**, HJAUP M0109).

Etymology: refers to the obovoid conidial shape.

Colonies on natural substrate effuse, brown to dark brown, hairy. Mycelium partly superficial, partly immersed in the substratum, composed of branched,



Fig. 1. *Hemicorynespora obovoidea* (holotype, HJAUP M0109): A–D. conidiophores and conidia; E, F. conidiophores; G, H. conidia.

septate, pale brown to brown, smooth hyphae. Conidiophores macronematous, mononematous, single, erect, unbranched, straight or flexuous, cylindrical, brown to dark brown, smooth, septate, 55–110 × 4–5.5 μm. Conidiogenous cells monotretic, integrated, terminal, cylindrical, brown, smooth, 16.5–34.5 × 4.5–5.5 μm, with 0–1 cylindrical percurrent extension. Conidial secession schizolytic. Conidia solitary, acrogenous, dry, obovoid, smooth-walled, brown, asymmetrically 1-septate, 16.5–20.5 × 8–11 μm, 1.5–2.5 μm wide at the truncate base.

Comments—Ellis (1972) established *Hemicorynespora* with two species, *H. deightonii* M.B. Ellis (type species) and *H. mitrata* (Penz. & Sacc.) M.B. Ellis. The genus is characterized by solitary, acrogenous, mitre-shaped, limoniform, obovoid or ellipsoidal, 0–1-septate, smooth conidia seceding schizolytically from integrated, terminal, monotretic, determinate or percurrently elongated conidiogenous cells (Ellis 1972, Seifert et al. 2011). Subsequently, 10 additional species have been reported in *Hemicorynespora* (Ellis 1972, Matsushima 1981, Holubová-Jechová 1987, Subramanian 1995, Mercado et al. 1997, Sivanesan &

Chang 1997, Delgado et al. 2007, Ma et al. 2012a). However, Ma et al. (2012a) excluded *H. obclavata* Subram. (with 2-septate conidia) and *H. multiseptata* Sivan. & H.S. Chang (with 4–5-septate conidia) from the genus; thus *Hemicorynespora* currently contains 10 recognized species. Only *H. clavata* G. Delgado et al. and *H. dimocarpi* Jian Ma & X.G. Zhang have previously been reported from China (Ma et al. 2012a).

Hemicorynespora obovoidea superficially resembles *H. ovata* Subram., which differs by its aseptate conidia (Subramanian 1995).

Solicorynespora jinggangshanensis Jian Ma & X.G. Zhang, **sp. nov.** Fig. 2

MycoBank MB 814892

Differs from *Solicorynespora melicopes* by its smaller, predominately 6-septate conidia with the basal cell longer than the second cell and from *S. obclavata* by its wider conidia with a shorter rostrum and the basal cell longer than the second cell.



Fig. 2. *Solicorynespora jinggangshanensis* (holotype, HJAUP M0290): A, B. conidiophores and conidia;. C. conidiophore with developing conidium; D, E. conidia.

Type: China, Jiangxi Province: Jinggangshan Mountain, on decaying twigs of an unidentified broadleaf tree, 6 Nov. 2014, J. Ma (**Holotype**, HJAUP M0290).

Etymology: refers to the locality where the type specimen was found.

Colonies on natural substrate effuse, brown to dark brown, hairy. Mycelium partly superficial, partly immersed in the substratum, composed of branched, septate, pale brown to brown, smooth-walled hyphae. Conidiophores macronematous, mononematous, single, erect, straight or flexuous, unbranched, smooth, septate, brown to dark brown, 60–145 × 5–6 μm. Conidiogenous cells monotretic, integrated, terminal, cylindrical or lageniform, brown to pale brown, smooth, 11.5–22.5 × 4.5–6.5 μm, with 0–3 percurrent proliferations. Conidial secession schizolytic. Conidia solitary, dry, acrogenous, obclavate, rostrate, smooth, brown, 5–7-euseptate (mostly 6-septate), 35.5–51 × 13.5–15.5 μm, 3.5–4.5 μm wide at the truncate base, apex extended into a subhyaline to pale brown rostrum, 2.5–3.5 μm wide.

Comments—Castañeda-Ruíz & Kendrick (1990), who re-examined the genus *Corynespora* Güssow using conidial septation as the circumscribing character, erected the segregate genus *Solicorynespora* to accommodate those species previously described in *Corynespora* but with solitary, euseptate phragmoconidia (McKenzie 2010, Ma et al. 2012b, Hernández-Restrepo et al. 2014). Thus far, 22 species have been described in or transferred to the genus (Hernández-Restrepo et al. 2014, Ma et al. 2014).

Of the known species, *Solicorynespora jinggangshanensis* is most similar to *S. melicopes* Jian Ma & X.G. Zhang and *S. obclavata* (Dyko & B. Sutton) R.F. Castañeda & W.B. Kendr. in conidial shape, but *S. melicopes* differs by its larger, predominately 5-septate, occasionally distoseptate conidia (77–94 × 19.5–27 μm); with the longest cell second from the base (Ma et al. 2012b); and *S. obclavata* differs by its narrower conidia (9.5–11 μm wide) with a longer rostrum and the longest cell second from the base (Castañeda-Ruíz & Kendrick 1990).

Solicorynespora lushanensis Jian Ma & X.G. Zhang, **sp. nov.** Fig. 3

MycoBank MB 814893

Differs from *Solicorynespora linderae* by its smooth, smaller conidia with more septa.

Type: China, Jiangxi Province: Lushan Mountain, on decaying twigs of unidentified broadleaf trees, 8 Nov. 2014, J. Ma (**Holotype**, HJAUP M0199).

Etymology: refers to the locality where the type specimen was found.

Colonies on natural substrate effuse, brown, hairy. Mycelium partly superficial, partly immersed in the substratum, composed of branched, septate, pale brown to brown, smooth-walled hyphae. Conidiophores macronematous,



Fig. 3. *Solicorynespora lushanensis* (holotype, HJAUP M0199): A. conidiophores; B, C. conidiophores and conidia; D. conidia.

mononematous, solitary or in groups, erect, straight or flexuous, unbranched, smooth, brown, 55–115 × 5–6 μm. Conidiogenous cells monotretic, integrated, terminal, cylindrical, brown, smooth, 9.5–20.5 × 4–5 μm, sometimes with 1 percurrent extension. Conidial secession schizolytic. Conidia solitary, dry, acrogenous, obclavate, tapering towards the apex, smooth, brown, paler towards the apex, 9–13-euseptate, 56–78 × 10–12 μm, tapering to 2–3 μm near the apex, 2.5–3.5 μm wide at the truncate base.

Additional specimen examined: **China**, **Jiangxi Province**: Lushan Mountain, on decaying twigs of unidentified broadleaf trees, 8 Nov. 2014, J. Ma (HJAUP M0190).

Comments—*Solicorynespora lushanensis* is similar in conidial shape to *S. linderae* Jian Ma & X.G. Zhang, which differs by its larger, 7–9 septate conidia (100–130 × 12.5–15.5 µm) with a verrucose basal cell and a pale brown, aseptate rostrum (Ma et al. 2012b).

Acknowledgments

The authors express gratitude to Dr. R.F. Castañeda-Ruíz and Dr. Eric H.C. McKenzie for serving as pre-submission reviewers and to Dr. Shaun Pennycook for nomenclatural review. This project was supported by the National Natural Science Foundation of China (No. 31360011) and the Education Department of Jiangxi Province of China (No. GJJ13282).

Literature cited

Castañeda-Ruíz RF, Kendrick B. 1990. Conidial fungi from Cuba: II. Univ. Waterloo Biol. Ser. 33. 61 p.

Delgado G, Mercado SA, Mena J, Guarro J. 2007. *Hemicorynespora clavata*, a new hyphomycete (anamorphic fungi) from Cuba. Crypt. Mycol. 28: 65–69.

Ellis MB. 1972. Dematiaceous hyphomycetes. XI. Mycol. Pap. 131. 25 p.

Hernández-Restrepo M, Castañeda-Ruíz RF, Gené J, Silvera-Simón C, Cano J, Guarro J. 2014. Two new species of *Solicorynespora* from Spain. Mycol. Progress 13: 157–164. http://dx.doi.org/10.1007/s11557-013-0903-9

Holubová-Jechová V. 1987. Studies on hyphomycetes from Cuba VI. New and rare species with tretic and phialidic conidiogenous cells. Česká Mykol. 41: 107–114.

Ma J, Ma LG, Zhang YD, Castañeda-Ruiz RF, Zhang XG. 2012a. New species or record of *Corynesporopsis* and *Hemicorynespora* from southern China. Nova Hedwigia 95: 233–241. http://dx.doi.org/10.1127/0029-5035/2012/0030

Ma J, Zhang YD, Ma LG, Ren SC, Castañeda-Ruíz RF, Zhang XG. 2012b. Three new species of *Solicorynespora* from Hainan, China. Mycol. Progress 11: 639–645. http://dx.doi.org/10.1007/s11557-011-0775-9

Ma J, Zhang XG, Castañeda-Ruíz RF. 2014. *Ceratosporium hainanensis* and *Solicorynespora obovoidea* spp. nov., and a first record of *Bactrodesmiastrum obscurum* from southern China. Mycotaxon 127: 135–143. http://dx.doi.org/10.5248/127.135

Matsushima T. 1981. Matsushima mycological memoirs 2. Published by the author, Kobe, Japan

McKenzie EHC. 2010. Three new phragmosporous hyphomycetes on *Ripogonum* from an 'ecological island' in New Zealand. Mycotaxon 111: 183–196. http://dx.doi.org/10.5248/111.183

Mercado SA, Heredia AG, Mena PJ. 1997. Tropical hyphomycetes of Mexico I. New species of *Hemicorynespora*, *Piricauda* and *Rhinocladium*. Mycotaxon 63: 155–167.

Seifert K, Morgan-Jones G, Gams W, Kendrick B. 2011. The genera of hyphomycetes. CBS Biodiversity Series 9. 997 p.

Sivanesan A, Chang HS. 1997. *Chaetosphaeria ampulliformis* sp. nov. associated with a *Hemicorynespora* anamorph, and a key to *Hemicorynespora* species. Mycol. Res. 101: 845–848. http://dx.doi.org/10.1017/S0953756297003493

Subramanian CV. 1995 ["1992/1993"]. *Agrabeeja kavakapriya* gen. et sp. nov. and additions to *Hemicorynespora*. Kavaka 20/21: 1–9.